

Fix the root cause and get back on your feet quickly

in a walking boot 1,2

"It did not take as long as I had anticipated to feel better and to live just as I had, if not better, than before."

April, Lapiplasty patient

**Explore** Life After

**Lapiplasty**®

3D Bunion Correction™

The #1 Bunion Product in the US<sup>3</sup>

**Bunions** are more than a bump

A common misconception is that a bunion is simply an overgrowth of bone that can be "shaved off". In reality, a bunion is a complex, 3D

deformity caused by an unstable joint in the middle of the foot which allows the metatarsal bone to drift out of visual "bump normal alignment.

actual root cause; an unstable joint

metatarsal bone >

Bunion foot



Did you know?

of adults in the U.S. have bunions4



Bunions can be mild, moderate, or extreme.

## Bunions are a progressive disorder and do not go away on their own without surgical correction5

Conservative treatments such as bunion pads/splints, toe spacers, orthotics, and wider toe-box shoes can help alleviate pain; but will not fix the problem.

However, Lapiplasty® 3D Bunion Correction™ can effectively treat bunions of all severities.



"My bunions were a genetic thing. They hurt. And, they were embarassing." Emily, Lapiplasty® patient

## Bunions are a 3D problem

While 87% of bunions are 3D problems with abnormal rotation of the metatarsal bone, the majority of bunion surgery is only 2D and fails to secure the root cause - an unstable joint.


When all three-dimensions of a bunion are not addressed, there is a 12X likelihood it will recur.7

That's why Lapiplasty® 3D Bunion Correction™ was developed; to give patients a better outcome.

metatarsal bone 🕨

Bunion foot

**Bunions are complex 3D deformities** that cause your big toe bones to:



**Elevate** Lean unstable foundation

First, like the Leaning Tower of Pisa, with its unstable foundation, the metatarsal bone starts to lean sideways.

Secondly, the metatarsal bone often elevates up, like a stool with a short leg; transferring excessive load to other toes.

Did you know?

of bunions are misaligned in 3 dimensions<sup>6</sup>





Thirdly, the metatarsal bone can rotate, causing abnormal wear and tear on your joints like a car tire out of alignment.

## Lapiplasty

3D Bunion Correction™


Fix it Right the First Time™

## **How it works**

New way

## **Lapiplasty® 3D Bunion Correction™**

Rotates bone back to normal 3D alignment. Reliably secures bunion's root cause.



- ✓ Returns entire bone to normal alignment; a 3D correction
- Fixes the root cause of the bunion; an unstable joint
- √ You'll be on your feet in a boot, in many cases, within 2 weeks¹,²


Did you know? Up to

of **2D** bunion surgeries (osteotomies) can recur over time<sup>8</sup>

Old way

## Traditional 2D Bunion Surgery (Osteotomy)

Shaves off "bump"; cuts & shifts top of bone. Does not address bunion's root cause.



- X Unnaturally cuts & shifts bone; only a 2D correction
- X Addresses cosmetic "bump" only; not the root cause
- X You may be off your feet for up to 6-8 weeks



"As an engineer, I was encouraged by the **precision** this technology provided. This option seemed like **common sense medicine** to me. I wanted the problem **fixed completely and correctly the first time** – who wouldn't?"

Tom, Lapiplasty® patient

# What should I expect in terms of recovery?

One Lapiplasty\* benefit, that our patients consider a "game changer" when it comes to recovery, is the ability to quickly bear weight in a boot.<sup>1,2</sup>

Although timepoints and doctor's orders vary based on each patient's specific condition, a typical recovery timeline is detailed below. Be sure to ask your doctor what your particular recovery protocol will look like to align your expectations.

## Within 2 weeks

You can begin to put some weight on your foot 1,2



Next 4-6 weeks

Walking in a boot



At 6-8 weeks

Move into comfortable shoes



At 4-6 months

Resume most activities & return to normal shoewear





## The results speak for themselves









Dr. Paul Dayton (IA), 7 weeks post-op

Dr. J P McAleer (MO), 12 weeks post-op









Dr. Daniel Hatch (CO), 4 months post-op

Dr. Alan Berman (NY), 2 years post-op










Example of pre- and post-op xrays

Example of pre- and post-op xrays

Visit AlignMyToe.com to see many more cases.

## Only the Lapiplasty<sup>®</sup> Procedure is backed by 15 publications and offers:

- 97-99% of patients maintain their 3D correction (as demonstrated in 13 &17 month studies)<sup>1,9</sup>
- <2 weeks to return to weight-bearing in a boot<sup>1,2</sup>
- 10.4mm average reduction in foot width<sup>10</sup>
- 2.4-3.1mm average shortening of the big toe<sup>11</sup>
- 3% of patients require hardware removal<sup>2</sup>



# You have questions We have answers ... from doctors\*

## Is the Lapiplasty\* Procedure covered by insurance and Medicare?

"Yes. Insurance and Medicare typically cover bunionectomy and joint fusion procedures that utilize the Lapiplasty® System when deemed medically necessary. Your doctor's office can reach out to your provider to determine your level of coverage. This process is best left to your doctor's office."

## Why was the Lapiplasty® Procedure developed?

"Lapiplasty" was born from doctors' dissatisfaction with bunion surgery. It's a philosophy of fixing the deformity in 3D and being able to do it in a reproducible way so that it's good for the doctor and patient."

## Can I have Lapiplasty\* 3D Bunion Correction™ if I've had previous 2D surgery that's failed?

"Yes. It is possible to have a failed 2D bunion surgery corrected and have an excellent clinical outcome."

## Will the titanium plates limit mobility?

"The plates will not limit functionality. By stabilizing and fusing the unstable joint, we are actually making the foot more functional."

#### Why do you offer the Lapiplasty® Procedure?

"Patients are often times scared of bunion surgery since they've heard horror stories from family members or neighbors – and sometimes they have experienced recurrence themselves. It's exciting to offer patients the Lapiplasty® Procedure with excellent outcomes, 8,9 good anatomic restoration, and an excellent recovery profile. It's a game changer."

#### Can I have both feet treated at the same time?

"It's possible, but not recommended. I like my patients to have one good mobile foot to rest/rely on during recovery. Walking in two cam boots is bulky and not recommended."

#### When can I return to sports / activities?

"Most patients can return to low-impact activities at 3 months after surgery; and impact sports at 4-6 months."

## When can I drive if my right foot is treated?

"Right around the 6-8 week mark, postoperatively, as I like my patients to be out of their walking boot."

## Are there physical limitations after the surgery?

"There are no functional limitations once a Lapiplasty® patient is fully healed."

#### What do your patients like about Lapiplasty\*?

"My patients like the results they are seeing when they look at their before & after xrays. It's the results, the early-weightbearing (in a boot), and the relief from pain."

#### Will my bunion come back?

"Whereas no surgical procedure has a 100% success rate, the Lapiplasty Procedure is designed with the intent to prevent recurrence. Data shows that 97 and 99% of patients were able to maintain their 3D correction in 13 and 17-month trials, respectively.<sup>8,9</sup> "

Visit **AlignMyToe.com** for many more FAQs.

<sup>\*</sup>Physicians are paid consultants of Treace Medical Concepts, Inc.

"When my doctor explained how the Lapiplasty" Procedure differed from other bunion treatments, I felt more at ease."

Carrie, Lapiplasty® patient



Visit AlignMyToe.com for more patient stories.

Did you know?

Lapiplasty® is the



bunion correction product in the U.S.<sup>3</sup>

Patient demand has made the Lapiplasty®
Procedure the #1 bunion product in the U.S.<sup>3</sup>

Of course, each individual's experience is unique to them and results vary, but we invite you to visit AlignMyToe.com to read more patient stories and to learn more about the procedure, including the risks associated with surgery.

Only a surgeon can tell if Lapiplasty® 3D Bunion Correction™ is right for you.

There are potential risks with surgery and recovery takes time. Potential risks include, but are not limited to: infection, discomfort from the presence of the implant, loosening of the implant, and loss of correction with nonunion or malunion - which means that the bone does not heal correctly.

For more information, including benefit and risk information about the procedure, please visit **www.AlignMyToe.com.** 



"The Lapiplasty Procedure enabled me to resume the things that **bring me joy.**"

Andrea, Lapiplasty® patient

#### References

- 1. Ray J, et al. Foot Ankle Int. 2019 Aug;40(8):955-960.
- 2. Dayton P, et al. J Foot Ankle Surg. 2019. 58:427-433.
- 3. Trailing 12mo. revenue based on company market estimates.
- 4. Kim Y, et al. Foot Ankle Int. 2015. 36:944-52.
- 5. American College of Foot & Ankle Surgeons website
- 6. Kim Y, et al. Foot Ankle Int. 2015. 36:944-52.
- 7. Okuda R, et al. JBJS, 2007, 89:2163-72.
- 8. Jeuken RM, et al. Foot Ankle Int. 2016. 37:687-95.
- 9. Dayton P, et al. J Foot Ankle Surg. 2020, 59(2): 291-297.
- 10. Vaida J, et al. Foot & Ankle Othopaedics. 2020. Vol. 5(3)1-5.
- 11. Hatch D, et al. Foot & Ankle Ortho. 2020, 5(4): 1-8.

# Lapiplasty® 3D Bunion Correction™

For more information, including benefit and risk information about the procedure, please visit www.AlignMyToe.com.



The Leader in Hallux Valgus Surgery™

Registered marks and trademarks of Treace Medical Concepts, Inc. Pat. treace.com/patents © 2021 Treace Medical Concepts, Inc. All Rights Reserved. M018E