## Q: What is the Percuplasty™ System?

**A:** A cutting-edge, minimally invasive surgical procedure to correct your bunion through small incisions on the side of your foot.




Individuals results may vary.

## Q: What are the benefits of a minimally invasive bunion procedure?

**A:** The key benefits:

- Smaller scars
- Less postoperative pain and swelling[1,2]
- Less need for opioids postoperatively[3]

# Percuplasty™

Percutaneous 3D Bunion Correction®

References
1. Lee M, et al. Foot Ankle Int. 2017 Aug;38(8):838-846.
2. Kheir E, et al. Foot Ankle Int. 2015 Mar;36(3):248-52.
3. Chan JJ, et al. Orthopedics. 2020 Jul 1;43(4):e334-e337.
4. Kim Y, et al. Foot Ankle Int. 2015. 36:944-52.
5. Okuda R, et al. JBJS. 2007. 89:2163-72.
6. Lai MC, et al. Foot Ankle Int. 2018 Mar;39(3):311-317.
7. Schipper O, et al. Orthop Clin N Am 51 (2020) 403–422.
8. Li G, et al. J Orthop Translat. 2024 Mar 11:45:48-55.

Individual results may vary depending on age, weight, health and other variables.

There are risks and recovery takes time.

Risks include infection, pain, implant loosening and loss of correction with improper bone healing. If any of these occur, additional treatments may be needed.


For more information on benefits, risks and recovery, visit **treace.com**.

*Based on typical 7mm incision size. Individuals results may vary.

TREACE®
*A Step Ahead.™*

Registered marks and trademarks of Treace Medical Concepts, Inc. Pat. treace.com/patents
®2025 Treace Medical Concepts, Inc. All Rights Reserved. M4864A

Percuplasty™
Percutaneous 3D Bunion Correction®




# Small Scar*, Huge Relief: Rapid Recovery After MIS Bunion Surgery Starts HERE.

# Bunions are a 3D problem

The Percuplasty™ System addresses more than just the “bump”. While 87% of bunions are 3-dimensional problems[4] with abnormal rotation of the metatarsal bone, the majority of bunion surgery is only 2-dimensional.

When all three dimensions of a bunion are not addressed, there is a 12X likelihood it will recur.[5]

**The Percuplasty™ System can help relieve discomfort through controlled 3D re-alignment of the metatarsal bone through small and discreet incisions.**




# MIS bunion surgery allows for smaller scars.[6]



**Percuplasty™ Percutaneous 3D Bunion Correction Surgery®**



**Traditional Bunion Surgery**

# Q: What could I expect during recovery after MIS bunion surgery?[7,8]



**Up to 6 Weeks**
of protected weight-bearing.



**At 6 Weeks**
May return to regular shoes and typically resume low impact activities.



**Around 3-4 Months**
Typically resume full activity.

Varies by patient as care differs depending on the individual’s needs and surgeon’s medical opinion/preference.